

Contents

- A. Afonin, B. Kopzhassarov, E. Milyutina, E. Kazakov, A. Sarbassova and A. Seisenova
Prototype Spatio-temporal Predictive System of pest development of the codling moth, *Cydia pomonella*, in Kazakhstan 1-12
- A. Sumbul and I. Mahmood,
Interactive effect of *Meloidogyne incognita* and *Macrophomina phaseolina* on the development of root-rot disease complex in relation to growth and physiological attributes of chickpea 13-23
- A. Archidona-Yuste, C. Cantalapiedra-Navarrete, J.E. Palomares-Rius, P. Castillo and E.A. Tzortzakakis
Plant-parasitic nematodes associated with cultivated and wild olive trees in Crete, Greece 24-28
- E.G. Badieritakis, A.A. Fantinou and N.G. Emmanouel
A qualitative and quantitative comparison of mite fauna between bifenthrin-treated and non-pesticide treated alfalfa hay fields in Central Greece 29-41
- A. Michaelakis, E. Anastasaki, P.G. Milonas, D.P. Papachristos, D. Kontodimas, C.M. Pontikakos, D.G. Raptopoulos, N.A. Babilis and M.A. Konstantopoulou
Efficacy of communication disruption of *Thaumetopoea pityocampa* (Lepidoptera: Thaumetopoeidae) with low pheromone formulation 42-53